

Autism is a range of complex neurobiological developmental disabilities which are defined by social interaction impairments, communication difficulties, and repetitive behaviors (Autism Speaks, 2016f). Autism occurs in all ethnic and socioeconomic groups. Approximately 1 in 68 children in the United States have some form of developmental delay known as an autism spectrum disorder (ASD), which is a higher estimate than the 1 in 150 estimates in 2000 and 2002 (Centers for Disease Control and Prevention [CDC], 2016). In acknowledging the significance of this dramatic increase from previous data, the Centers for Disease Control and Prevention (CDC) refers to autism as an “important public health concern” and thus supports extensive research to understand why autism occurs (CDC, 2016).

#### **AUTISM SPECTRUM DISORDERS (ASD)**

After a 14-year review process, the American Psychiatric Association published the fifth edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5) in May 2013. The DSM-5 is used by physicians to diagnose and classify mental disorders, including Autism. The diagnoses criteria for Autism were updated in the DSM-5. The change was recommended as a better reflection of the state of knowledge of Autism. The new single umbrella disorder, Autism Spectrum Disorder, will encompass the separate diagnosis labels of Asperger’s Syndrome, Autistic Disorder and Pervasive Development Disorder. The change is expected to improve the diagnosis of ASD and should not substantially change the number of children being diagnosed (American Psychiatric Association, 2016). Current published materials on Autism

reflect the terminology of the DSM-4 and the information that follows explains those terms.

Autism is one of three disorders classified in the ASD. The three disorders have similar symptoms and signs in problems with social interaction, communication, and behaviors, but differ in severity, occurrence, and the nature of symptoms (CDC, 2016b).

A pervasive developmental disorder (PDD) is another term commonly used to categorize autism. PDD is an umbrella term under which the specific diagnoses are defined. All three disorders in ASD are considered to be part of PDD. Some confusion and debate has occurred among professionals in the autism field on how to correctly refer to and categorize autism, but generally autism literature will refer to autism as being an ASD and/or a PDD (Autism Speaks, 2016d). It may be some time before the medical journals and professional literature reflect the DSM-5 terminology for Autism (ASD) exclusively.

- **Autistic Disorder** – Individuals with this disorder experience difficulties in social interaction, communication, and display unusual behaviors and interest. They suffer from delays in cognitive abilities and language disorders; it is commonly referred to as “classic” autism (CDC, 2016b).
- **Asperger Syndrome** – Individuals who suffer from this disorder have difficulties in social interaction and show a limited range of interests and/or repetitive behaviors. One difference between this disorder and autistic disorder is that persons affected with Asperger syndrome do not have

trouble with language, and generally have an IQ in the normal to superior range (Autism Speaks, 2016a).

- **Pervasive Developmental Disorder-Not Otherwise Specified** – Also called “subthreshold autism,” this group is for individuals who have autistic symptoms and signs, but who do not meet the specific diagnostic criteria of the other two disorders or have mild symptoms. For example, individuals may have extensive symptoms in one area, but moderate or no symptoms in another (Autism Speaks, 2016d).

**PREVALENCE OF AUTISM**

While current figures show that autism occurs in all facets of society, there are three groups that appear to have a higher-than-normal risk for autism:

- **Boys** – Males are almost five times more likely to have autism than females.
- **Siblings of those with a form of autism** – Families that have one child with autism show a rate of recurrence of autism in another child of about 2-18 percent (significantly higher than the general population).
- **Individuals with other diagnoses**– Autism occurs more often (around 10%) in people who have certain genetic and chromosomal disorders such as fragile X syndrome, Down syndrome, and tuberous sclerosis (CDC, 2016a).

*Autism Prevalence in Texas*

Children with autism, ages 3-21, constitute the fourth largest component of special education eligibility in Texas (Office of Special Needs, n.d.). The actual prevalence of autism is not entirely known, but an analysis of educational data provides perhaps the closest indications. The following table, with data from the Texas Education Agency (2016), illustrates the growth of autism cases in Texas among 3-21 year old students during the 2013-2015 school years. For the 2015 school year, students with autism represented 12% of the total students with disabilities population, up from 10.6% in 2013 and 11.4% in 2014.

**NUMBER OF CHILDREN WITH AUTISM IN TEXAS SCHOOL SYSTEM**

	2013	2014	2015
<b>Ages 3-5</b>	7,064	7,464	8,030
<b>Ages 6-21</b>	39,826	43,866	47,570
<b>TOTAL</b>	46,890	51,330	55,600

(Texas Education Agency, 2016)

*Autism Prevalence in San Antonio area*

According to data from the Texas Education Agency, the number of individuals ages 3-21 with an autism diagnosis and who are being educated in the Region 20 school system has increased during the past three school years. This is consistent with national studies showing that autism in children is increasing.

In Region 20, the numbers of autistic children are increasing at a much higher rate than the numbers of children with disabilities (of which autism is considered one). From 2013 through 2015 the number of autistic children in school increased 19% while all students with disabilities increased only 6%. The Region 20 school system includes school districts in the following counties: Atascosa, Bandera, Bexar, Dimmit, Frio, part of Guadalupe and Karnes, the majority of Kendall, Kerr, Kinney, La Salle, Maverick, Medina, Real, Uvalde, Wilson, and Zavala (Educational Service Center, Region 20; 2016). This data represents the most recent figures released.

**NUMBER OF CHILDREN WITH AUTISM IN REGION 20 SCHOOL DISTRICT**

	2013	2014	2015
<b>Disability</b>	41,569	42,824	43,913
<b>Autism</b>	4,130	4,445	4,905

(J. Enriquez, Personal Communication, October 3, 2016)

**DIAGNOSING AUTISM**

Researchers have determined that early diagnosis and intervention has a noticeable impact on an autistic child’s development. It is recommended that children be tested for autism at 9, 18 and 24 months of age (CDC, 2016e).

The signs and symptoms of autism appear in varying degrees and are classified under three main categories: social skill impairments, communication impairments and repetitive behaviors.

*Social Deficits*

- Impaired use of nonverbal behaviors (eye contact, facial expression, gestures, etc.) to regulate social interaction
- Little seeking to share interests with people, lack of development of peer relationships
- Lack of social-emotional reciprocity

*Communication Deficits*

- Delay in or absence of spoken language
- Distinctive or repetitive use of language
- Limited ability for make-believe play
- Inability to maintain a conversation with others

*Repetitive and Stereotyped Behavior*

- Preoccupations with one or more stereotyped interests that is abnormal in intensity or focus
- Adherence to specific rituals
- Stereotyped body movements (hand or finger flapping or twisting, etc.)
- Persistent preoccupation with parts or sensory qualities of objects

(Autism Speaks, 2016e)

While some parents may be hesitant in taking their children to see a physician and initially attribute the symptoms as temporary, the following “red flags” should signal for immediate evaluation by a qualified physician:

**“RED FLAGS”**

- ✓ Does not offer joyful expressions, such as big smiles, by six months of age
- ✓ Does not reciprocate facial expressions by the age of nine months
- ✓ Does not babble by 12 months
- ✓ Does not gesture (point, wave, etc.) by 12 months
- ✓ No words by the age of 16 months
- ✓ Does not say two-word phrases (excluding repeating and/or imitating) by 24 months
- ✓ Has any loss of language or social skills at any age

(Autism Speaks, 2016b)

Currently, no medical tests are available to detect autism. However, Australian researchers have reported that they have developed a genetic test that can predict autism with those of central European descent with a 70% accuracy rate, which can lead to earlier and more precise detection in the future (Heasley, 2012).

**POSSIBLE CAUSES**

While there are various projects investigating possible causes of autism, there are no definite known reasons for its occurrence at this time. A variety of factors linked to autism continue to be investigated. These factors include genetics, the environment, neurobiology, and the mother’s pregnancy.

*Genetics*

Evidence supporting the link between genetics and autism can be illustrated by examining the frequency of twins with autism. A recent study found that when autism was identified in one twin the other twin had a high incidence of having autism. The concordance rate for fraternal twins has been shown to be between 0-31%, while the rate among identical twins is estimated to be approximately 36-95% (CDC, 2016a).

A study published in September 2015 identified 239 genes which, if disabled by a mutation, had a high likelihood of causing autism. It found that “in at least half of cases, autism traces to one of roughly 200 gene-disabling mutations found in the child but neither parent. Many of these “high-impact” mutations... completely disable genes crucial to early brain development” (Autism Speaks, 2015). In 2016, one study reported that mutations in non-coding regions of the genome in a parent’s eggs or sperm can have a major effect on early brain development contributing to autism and that “most autism-linked de novo mutations come from the father and tend to increase with his age” (Autism Speaks, 2016c). Other evidence supporting genetic causes of autism includes genetic vulnerability. Autism tends to occur more frequently when certain other conditions are present including; Fragile X syndrome, tuberous sclerosis, congenital rubella syndrome, and untreated phenylketonuria (Autism Society of America, n.d.a).

*The Environment*

A study published in the Archives of General Psychiatry found that 40% of autism risk was linked to genetics, while 55% was connected to environmental factors (Pearson, 2011). It has been found that many children with autism have a

metabolic impairment that reduces their ability to rid the body of heavy metals and toxins, leading to brain/nervous system damage and developmental delays (Autism Society of America, n.d.b). A study conducted at the UT Health Science Center in San Antonio showed a statistically significant link between industrial release of mercury and autism. The researchers discovered that the prevalence of autism was reduced by 1 to 2 percent for every 10 mile distance from the pollution source (UT Health Science Center San Antonio, 2008). More recently, a study published in 2015 reported that there was a significant increase in autism prevalence in children residing within 28 miles of air pollutant releasing industrial facilities (Dickerson, et al., 2015).

In prior years, it was believed that the increase of autism cases was linked to children's vaccines. This belief has been entirely disproved. The CDC (2016c) states, "studies continue to show that vaccines are **not** associated with ASDs."

### *Neurobiology*

The emergence of brain imaging technologies such as CT, MRI, and PET scans has allowed researchers to identify numerous areas of the brain involved in autism. Also being studied are the roles of neurotransmitters such as serotonin, dopamine, and epinephrine. It has been shown that abnormal brain development in a child's first months may be a contributing cause of autism (Brain & Behavior Research Foundation, 2016). In a review of current research, researchers suggest that injury to the cerebellum (which contains nearly half of the brain's nerve cell connections) either before or during birth results in a significantly increased risk of autism (Autism Speaks, 2014b).

### *Pregnancy*

Metabolic conditions (diabetes, hypertension, and obesity) during pregnancy may be associated with ASD (Krakowiak et al., 2012). A new study by researchers at the University of California-Davis MIND Institute showed that high daily iron intake reduced the risk of having an Autistic child by 50% compared to mothers

not taking iron supplements (Autism Speaks, 2014a). The age of the mother is another factor; the older the mother during pregnancy, the higher risk of autism (Sandin et al., 2012). Prenatal exposure to valproic acid and thalidomide has been linked to higher risks of ASD (CDC, 2016b).

### **TREATMENT**

No known cure for autism exists nor does one single treatment package help all individuals with autism. Doctors specializing in diagnosing ASD will evaluate the child's condition and often provide a variety of different treatments that will help improve his/her lifestyle. The available treatments include behavioral, educational and medical interventions.

#### *Behavioral Interventions*

Applied behavior analysis (ABA) is an umbrella term used to describe interventions that work to reinforce desired behaviors and reduce unwanted behaviors. Though different in implementation methods, the Lovaas Model and the Early Start Denver Model (ESDM) both are intensive instruction models that utilize the techniques of ABA for young children. Social skills and play/interaction interventions can be provided through speech-language and occupational therapists. Cognitive Behavioral Therapy (CBT) can also be implemented to assist with behavioral challenges (Agency for Healthcare Research and Quality [AHRQ], 2014).

#### *Educational Interventions*

Educational interventions are the primary treatment for ASD and are intended to be administered in an educational setting. An educational intervention program frequently used is the Training and Education of Autistic and Related Communication Handicapped Children (TEACCH), which focuses on individualized assessments and emphasizes the preference of Autistic individuals for visual information processing (Autism Speaks, n.d.). The 81<sup>st</sup> Texas Legislature (2009) created the Texas Autism Research and Resource Center (TARRC) in response to the needs of the autism community. TARRC provides support to families, coordination of services by local groups, training and education for

professionals, current research on autism, and information on university research programs in Texas related to ASD (TARRC, 2016).

### *Medical Interventions*

Medication is often only prescribed to alleviate some symptoms associated with autism.

Among drugs used are selective serotonin reuptake inhibitors (SSRIs), anti-psychotics, stimulants, and anti-anxiety drugs (AHRQ, 2014). Some have seen an improvement in autistic symptoms with the usage of dietary supplements and/or a gluten-free, casein-free diet (GFCF); however at this time there is not enough research to prove that diet change has an impact on autism (AutismWeb, 2016).

### **COST**

Caring for a child with Autism is expensive. “It is estimated to cost at least \$17,000 more per year to care for a child with ASD compared to a child without ASD. Costs include health care, education, ASD-related therapy, family-coordinated services, and caregiver time. For a child with more severe ASD, costs per year increase to over \$21,000” (Supported Academics & Independent Life Skills, 2014). Additionally, some intensive behavioral therapy interventions can cost \$40,000 to \$60,000 per year per child (CDC, 2016a).

### **AUTISM RESOURCES IN SAN ANTONIO**

Any Baby Can (ABC) of San Antonio’s “Autism Services Program” includes parent education services for families of children who have been diagnosed with ASD, several autism support and education groups including one in Spanish, and group and individual counseling with a licensed psychologist. These services are provided at ABC’s Bexar, Comal and the 17 surrounding county locations. There also is an autism library with books, videos, and magazines, and an autism resource guide that provides additional information about autism services in the San Antonio area. In addition to monthly events, there are “Parent’s Night Out” evenings, which provide parents with some autism-free “me” time (Any Baby Can San Antonio, 2013).

The Early Childhood Intervention (ECI) program is a statewide program administered by the

Texas Department of Assistive and Rehabilitative Services (DARS) that focuses exclusively on services for infants from birth to age three that suffer from disabilities or developmental delays, including autism. The benefits offered by ECI include promoting development and learning, providing support to families, coordinating services, and decreasing the need for costly special programs (Texas Department of Assistive and Rehabilitative Services [DARS], n.d.a). There are four organizations in the San Antonio area that offer ECI services: the Brighton Center, Center for Health Care Services (CHCS), the Easter Seal Rehabilitation Center, and the Hill Country MH and DD Centers (DARS, n.d.b).

Autism Treatment Center, headquartered in Dallas, has a day program and several community-based group homes in San Antonio. Services provided include educational opportunities for persons ages 3-21, a diagnostic program which includes assessments and developmental autism screenings, and a Therapy Clinic that offers occupational and speech therapy (Autism Treatment Center, n.d.).

Founded in 2008, the Autism Service Center of San Antonio d.b.a. Autism Community Network was created with the long-term vision that it would become the “hub” for collaboration of autism services in the San Antonio area. Its mission to maximize the potential of children with autism, provide services to underserved children and families, and promote awareness in the community is encompassed in three primary goals to provide:

- Education and training to autism service providers
- Interdisciplinary diagnosis and treatment to young children with ASD
- Comprehensive information and networking for San Antonio’s autism community

(Autism Community Network, 2016)

Autism Lifeline Links (ALL) is a coalition of agencies and organizations founded through an initiative funded by the Kronkosky Charitable Foundation. ALL works as a one-time stop for families that are looking for autism related services ranging from diagnosis, respite care to

where an individual with autism can get a haircut. It includes nine (9) referral agencies who work directly with individuals with autism and 12 community partners including other local nonprofits and Universities.

ALL's referral agencies are part of a HIPPA-secured information sharing platform operated by TAVConnect. The platform allows only the referral agencies to share the information and it is fully designated to streamline and simplify connecting individuals and families with agencies and organizations that provide services addressing the challenges of Autism Spectrum Disorder (Autism Lifeline Links, 2016).

## REFERENCES

- Agency for Healthcare Research and Quality (AHRQ). (2014). *Therapies for children with Autism Spectrum Disorder*. Retrieved from <https://www.effectivehealthcare.ahrq.gov/search-for-guides-reviews-and-reports/?pageaction=displayproduct&productID=1974#9156>
- American Psychiatric Association. (2016). *What is Autism Spectrum Disorder?* Retrieved from <https://www.psychiatry.org/patients-families/autism/what-is-autism-spectrum-disorder>
- Any Baby Can of San Antonio. (2013). *Autism services*. Retrieved from <http://www.anybabycansa.org/services/autism-services/>
- Autism Community Network. (2016). *About us*. Retrieved from <http://www.acn-sa.org/index.cfm?fuseaction=Page.ViewPage&pageId=519>
- Autism Lifeline Links (2016). *About us*. Retrieved from [http://autismlifelinelinks.org/About-Us#tab\\_who](http://autismlifelinelinks.org/About-Us#tab_who)
- Autism Society of America. (n.d.a). *Differential Diagnosis*. Retrieved October 3, 2016, from <http://www.autism-society.org/what-is/diagnosis/differential-diagnosis/>
- Autism Society of America. (n.d.b). *Environmental Health* Retrieved October 3, 2016, from <http://www.as-az.org/environmental-health/>
- Autism Speaks. (n.d.). *Teaching children with Autism: Special education interventions*. Retrieved October 4, 2016, from [https://www.autismspeaks.org/sites/default/files/documents/family-services/faq\\_teaching.pdf](https://www.autismspeaks.org/sites/default/files/documents/family-services/faq_teaching.pdf)
- Autism Speaks. (2014a). *Taking iron during pregnancy & breastfeeding may lower Autism risk*. Retrieved from <https://www.autismspeaks.org/science/science-news/taking-iron-during-pregnancy-breastfeeding-may-lower-autism-risk>
- Autism Speaks. (2014b). *Studies implicate early Injury to cerebellum as major cause of autism*. Retrieved from <https://www.autismspeaks.org/science/science-news/studies-implicate-early-injury-cerebellum-major-cause-autism>
- Autism Speaks. (2015). *Study: Half of all autism cases trace to rare gene-disabling mutations*. Retrieved from <https://www.autismspeaks.org/science/science-news/study-half-all-autism-cases-trace-rare-gene-disabling-mutations>
- Autism Speaks. (2016a). *Asperger syndrome*. Retrieved from <https://www.autismspeaks.org/what-autism/asperger-syndrome>
- Autism Speaks. (2016b). *Learn the signs of autism*. Retrieved from <https://www.autismspeaks.org/what-autism/learn-signs>
- Autism Speaks. (2016c). *MSSNG Study expands understanding of Autism's complex genetics*. Retrieved from <https://www.autismspeaks.org/science/science-news/mssng-study-expands-understanding-autism%E2%80%99s-complex-genetics>
- Autism Speaks. (2016d). *PDD-NOS*. Retrieved from <https://www.autismspeaks.org/what-autism/pdd-nos>
- Autism Speaks. (2016e). *Symptoms*. Retrieved from <https://www.autismspeaks.org/what-autism/symptoms>
- Autism Speaks. (2016f). *What is autism?* Retrieved from <https://www.autismspeaks.org/what-autism>
- Autism Treatment Center (ATC). (n.d.). *San Antonio program*. Retrieved October 3, 2016, from <http://atcoftexas.org/sanantonio.html>
- AutismWeb. (2016). *The GFCF (gluten-free, Casein-free) diet for Autism spectrum disorders*. Retrieved from <http://www.autismweb.com/diet.htm>
- Brain & Behavior Research Foundation. (2016). *Frequently asked questions about Autism*. Retrieved from <https://bbrfoundation.org/frequently-asked-questions-about-autism>
- Centers for Disease Control and Prevention (CDC). (2016a). *Autism spectrum disorder (ASD): Data and statistics*. Retrieved from <http://www.cdc.gov/ncbddd/autism/data.html>
- Centers for Disease Control and Prevention (CDC). (2016b). *Autism spectrum disorder (ASD): Facts about ASD*. Retrieved from <http://www.cdc.gov/ncbddd/autism/facts.html>
- Centers for Disease Control and Prevention (CDC). (2016c). *Autism spectrum disorder (ASD): Related topics*. Retrieved from <http://www.cdc.gov/ncbddd/autism/topics.html>
- Centers for Disease Control and Prevention (CDC). (2016d). *Autism spectrum disorder (ASD): Research*. Retrieved from <http://www.cdc.gov/ncbddd/autism/research.html>
- Centers for Disease Control and Prevention (CDC). (2016e). *Autism spectrum disorder (ASD): Screening and diagnosis for healthcare providers*. Retrieved from <http://www.cdc.gov/ncbddd/autism/hcp-screening.html>
- Dickerson, A., Rahbar, M., Han, I., Bakian, A., Bilder, D., Harrington, R., ...Baio, J. (2015). *Autism spectrum disorder prevalence and proximity to industrial facilities releasing arsenic, lead or mercury*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/26218563>
- Educational Service Center, Region 20. (2016). *ESC-20 Districts*. Retrieved from <http://www.esc20.net/default.aspx?name=esc.R20.public.isds>
- Hesley, S. (2012). *Scientists develop genetic test for autism*. Retrieved from <http://www.disabilitycoop.com/2012/09/13/scientists-genetic-test/16445/>
- Krakiwaki, P., Walker, C., Bremer, A., Baker, A., Ozonoff, S., Hansen, R., & Hertz-Picciotto, I. (2012). *Maternal metabolic conditions and risk for autism and other neurodevelopmental disorders*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/22492772>
- Office of Special Needs. (n.d.). *Education directory for children with special needs: Texas*. Retrieved October 5, 2016, from [http://apps.militaryonesource.mil/MOS/?p=EFMP\\_DIRECTORY:HOME:0:::EKMT:37.20.10.180.0.0.0.0](http://apps.militaryonesource.mil/MOS/?p=EFMP_DIRECTORY:HOME:0:::EKMT:37.20.10.180.0.0.0.0)
- Pearson, C. (2011). *Autism: Environmental factors messing with our kids? (STUDY)*. Retrieved from [http://www.huffingtonpost.com/2011/07/04/studies-suggest-environment-autism-causes\\_n\\_889041.html](http://www.huffingtonpost.com/2011/07/04/studies-suggest-environment-autism-causes_n_889041.html)
- Sandin, S., Hultman, C., Kolevzon A., Gross, R., MacCabe, J., & Reichenberg, A. (2012). *Advancing maternal age is associated with increasing risk for autism: a review and meta-analysis*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/22525954>
- Supported Academics & Independent Life Skills. (2014). *Interesting autism spectrum disorder (ASD) statistics*. Retrieved from <http://www.sailsforstudents.org/autism-spectrum-disorder-statistics.htm>
- Texas Autism Research and Resource Center (TARRC). (2016). *About TARRC*. Retrieved from <http://legacy.hhsc.state.tx.us/tarrc/about/about-home.html>
- Texas Department of Assistive and Rehabilitative Services (DARS). (n.d.a). *Early childhood intervention services: What is ECI?* Retrieved October 3, 2016, from <http://www.dars.state.tx.us/ecis/index.shtml>
- Texas Department of Assistive and Rehabilitative Services (DARS). (n.d.b). *ECI program search*. Retrieved October 3, 2016, from <https://dmzweb.dars.state.tx.us/prd/citysearch>
- Texas Education Agency. (2016). *Demographic data—disability*. Retrieved from <http://www.cdc.gov/ncbddd/autism/facts.html>
- UT Health Science Center San Antonio. (2008). *Study links autism risk to distance from power plants, other mercury-releasing sources*. Retrieved from <http://www.uthscsa.edu/hscnews/singleformat2.asp?newID=2732>