

The population of Americans aged 65 and older numbered 49.2 million in 2018 (U.S. Census Bureau, 2020a) and is projected to reach 94.7 million in 2060. By 2040, there will be over 80 million older persons, over twice their number in 2010 (U.S. Census Bureau, 2020e).

The older population itself is increasingly older as well. Persons reaching age 65 have an average life expectancy of an additional 19.3 years; a child born in 2017 could expect to live 78.6 years, approximately 30 years longer than a child born in 1900 (U.S. Administration on Community Living [ACL], 2018b, p.2). The 85+ population is projected to more than double from 6.4 million in 2016 to 14.4 million in 2040 (U.S. Census Bureau, 2020e). These significant changes are not only relevant to those that are 65 years of age and older, but also to the remainder of society.

People 65 years and older make up 15.2% of the United States population, or one in every seven people; in Texas, they make up 12% of the population (U.S. Census Bureau, 2020a). The following table shows the percentage of persons per county in the San Antonio area that are 65 and older:

COUNTY	% OF POPULATION 65 AND OLDER IN 2018
Bandera	25.9%
Bexar	12.0%
Comal	17.7%
Kendall	18.8%
(U.S. Census Bureau, 2020a)	

This inevitable shift to an older population has a number of implications, and indicates a wider

spectrum of change already ongoing in American culture such as:

- changes in the perception of aging
- the overall quality of life for the elderly
- the practice of elder care
- the role of medicine for the elderly

With healthcare and technology in a constant state of advancement, the elderly population in the U.S. will continue to grow at high rates.

In Texas, “the 65 plus age category had the greatest increase (slightly more than one million) and grew at the fastest rate compared to the younger age groups” (Texas Demographic Center, 2019, p.1).

Additionally, the elder population itself is aging. In 2017 the 65-74-year-old population was over 13 times larger than in 1900, the 75-84 range was 19 times larger and the 85 and older category was 52 times larger. There were also 86,248 persons aged 100 or older in 2017, more than twice the 32,194 from 1980 (ACL, 2018b).

The elder population in 2018 included:

POPULATION 65 AND OLDER, 2018		
	U.S.	TEXAS
65-74 years	28,535,419	2,000,715
75-84 years	14,197,679	971,168
85+	6,205,483	365,931
Total 65+	49,238,581	3,337,814
(U.S. Census Bureau, 2020a)		

THE AGING PROCESS

Aging is a process of gradual maturation. Senescence is the biological process by which the capacity for cell division and the capacity for growth and function are lost over time, ultimately leading to death (Goldsmith, 2020).

Changes that occur during aging can affect individuals differently – some may develop diseases and impairments commonly associated with aging and others may not.

During the aging process a number of physiological changes begin to take place that not only affect appearance but also how the body functions and responds to daily living. Overall, the changes in most elderly involve a slowing down of all organ systems due to a decline in cellular activity (National Institute on Aging [NIA], n.d.). Normal physiological changes include:

Body Shape- Body fat increases up to 30% toward the center of the body (including the abdominal organs) as muscle and bone mass decreases. Changes in bones, muscles, and joints result in the tendency to become shorter, typically 1 cm (0.4 inches) for every 10 years after age 40. After turning 70, height loss occurs faster and an individual may lose a total of 1 to 3 inches in height (National Institutes of Health [NIH], 2020a).

Skin- Among the most visible signs of aging are wrinkles, sagging skin and graying hair. The skin gets thinner and more fragile and the layer of fat beneath the skin also thins. The result is a proneness to skin injury, and more than 90% of the elderly population has some kind of skin disorder (NIH, 2020e).

Senses- Taste, smell, touch, hearing, and vision all degrade with age. Nearly one in three of those aged 65 to 74 and 50 percent of those who are 75 and older have difficulty hearing (National Institute on Deafness and other Communication Disorders [NIDCD], 2018).

While almost all adults over 60 experience some visual loss, 6.2% of adults age 65+ in the state of Texas reported a visual disability in 2018 (U.S. Census Bureau, 2020b).

Problems with taste also increase with age. 19% of adults over 40 report some alteration in their sense of taste increasing to 27% of adults aged 80 and older (NIDCD, 2019).

Alterations in the sense of smell also occur with age. 23% of people over 40 and 32% of those

over 80 experience some smell dysfunction (NIDCD, 2019).

Taste bud decreases can result in a lack of interest in eating, while the diminishing sense of smell can lead to less awareness of personal hygiene. Touch sensation changes with age, which increases risk of frostbite, hypothermia, and burns. Reduced ability to perceive where the body is in relation to the floor increases the risk of falling (NIH, 2020d).

Nervous System- Normal changes include loss of nerve cells and weight in the brain and spinal cord as well as waste product build-up that causes abnormal structures to form. Normal decline of mental functions includes slowing of thought, memory and thinking (NIH, 2020c).

Cardiovascular System- Reduced blood flow through the body due to normal atrophy of the heart muscle, calcification of the heart valves, loss of elasticity in artery walls (arteriosclerosis) and intra-artery deposits (atherosclerosis) leads to a slower rate of healing, lower response to stress and increased risk of drug toxicity in addition to risk of hypertension, stroke, heart attack, or congestive heart failure (NIA, 2018).

Skeletal System- Loss of bone calcium can result in osteoporosis with increased risk of fracture. Arthritis is the degenerative inflammation of the joints and is the most common chronic condition within the elderly population. Osteoarthritis (joint cartilage wear) is the most common type of arthritis, and can inhibit mobility and daily activities (NIH, 2020b).

Muscular System- Loss of muscle tone and strength often manifests as a reduced ability to breathe deeply and/or reduced gastrointestinal activity, which can lead to constipation and bladder incontinence, particularly in women (Besdine, 2019).

Digestive System- While less affected by aging than other organ systems, certain digestive problems are more likely to develop in older adults, including conditions related to a decrease in muscular strength and elasticity: gastroesophageal reflux (GERD); difficulty swallowing; and constipation. Lactose

intolerance results from decrease in enzyme levels and excessive growth of bacteria leads to pain, bloating, and weight loss as well as decreased absorption of vitamin B12, iron, and calcium (Ruiz, 2019).

Endocrine System- Hormone levels and activity decrease leading to decreased muscle mass and increased risk of dehydration and type 2 diabetes (Besdine, 2019).

PHYSICAL HEALTH

The U.S. Department of Health and Human Services (HHS) (2016a) noted that while Americans could expect a longer life, they could also expect to develop any one of a number of chronic diseases and conditions associated with aging. Approximately two thirds of older adults have multiple chronic conditions. Although the mortality rates from heart disease, stroke, and cancer have been declining for several years, the incidence of those diseases is directly proportionate to age. The following table shows the percentage of selected chronic diseases for various age groups (most recent data available):

PERCENT OF POPULATION WITH CHRONIC DISEASE (2018)			
Chronic Disease	Age 45-64	Age 65-74	Age 75+
Hypertension	34.4	54.4	61.1
Arthritis	30.3	48.3	53.7
Hearing Trouble	17.8	31.6	47.2
Heart Disease	6.0	15.5	23.9
Cancer	9.6	22.2	31.3
Diabetes	12.9	22.2	22.8
Vision Trouble	15.9	16.9	19.5
Stroke	3.1	6.9	11.8
Kidney Disease	2.4	5.4	8.0
Emphysema	1.6	4.1	4.5

(CDC, 2019b)

The difference between the 45-64 age group and the older age groups signifies the higher prevalence of chronic disease that occurs in older people.

In general, the percentage of adults with good health declined with age. Those with higher education reported better health as did those with higher income and those living in a Metropolitan Statistical Area. Among adults aged 65 and over, 52.6% of those who had

Medicare and Medicaid had fair or poor health compared with 24.9% of those with only Medicare and 16.9% of those with private health insurance (CDC, 2019b).

Nutrition

As people age, food insecurity and hunger increase health risks. Research has demonstrated that seniors with low food security experienced more depression, lower quality of life, and reduced physical performance. In fact, the higher the level of food insecurity, the more issues with general health/functioning, pain, and mental health. Seniors are also more likely to have nutrient deficiencies and obesity (National Council on Aging [NCOA], n.d.b).

Reports show that the food insecurity rate for all senior households was 7.8% in 2016, up from 5.5% in 2001. “At the same time, the percentage of seniors facing the threat of hunger has more than doubled” (NCOA, n.d.b).

The National Council on Aging (n.d.a) reported that “16% of independent older adults are at high risk for malnutrition” and “up to 60% of older adults in health care settings are malnourished.”

One study found that a large portion of older adults do not get adequate amounts of nutrition from their diet and even 20% to 50% of those who used nutritional supplements still fell significantly short of the recommended amounts of folate, vitamin E, and magnesium (Sebastian, Cleveland, Goldman, Moshfegh, 2007). Vitamin D deficiency is the highest in the elderly, especially in those who have limited sun exposure (Kweder and Eidi, 2018).

MENTAL HEALTH

For at least the last decade, mental health has been recognized to be as important as physical health to the well-being of older Americans. It has been estimated that “over 20% of adults aged 60 and over suffer from a mental or neurological disorder” and “6.6% of all disability among over 60s is attributed to neurological and mental disorders” (World Health Organization [WHO], 2017).

Social Isolation

People need people. “Social connection influences a variety of mental and physical health outcomes. For example, those who are isolated are at increased risk for depression, cognitive decline, and dementia” (Holt-Lunstad, 2018).

Additionally, studies have shown that “poor social relationships were associated with a 29 percent increase in risk of coronary heart disease and a 32 percent rise in the risk of stroke” (Health Resource & Services Administration, 2019).

Social support includes emotional support such as sharing problems or venting emotions as well as soliciting advice and guidance. Other aspects include physical assistance such as transportation for shopping or doctor visits or even help with household chores. Seniors who live alone are more likely to succumb to depression as a result of social isolation.

In 2018, 29% of all households in the United States consisted of one or more individuals aged 65 or older. 37% of senior households consisted of individuals aged 65 and over living alone (U.S. Census Bureau, 2020c).

In its Senior Report 2018, America’s Health Rankings (2018) measured six factors to assess risk of social isolation:

- *Divorce, separated, or widowed:* living with a life partner decreases loneliness
- *Never married:* less likely to be socially integrated
- *Poverty:* lower income and less education leads to likelihood of social isolation
- *Disability:* physical or mental limitations are more likely to restrict social interaction
- *Independent living difficulty:* limited mobility increases isolation
- *Living alone:* “more likely to be poorly socially integrated and experience feelings of loneliness” (p.13)

(America’s Health Rankings, 2018)

Texas ranked 38th (of 50 states) for overall risk of senior social isolation but ranked in the bottom 10 for Disability and Poverty measures (America’s Health Rankings, 2018).

The table at the end of this brief shows the most recently available Household

Characteristics of seniors 65 and over in the U.S., Texas, and KCF counties of interest.

Depression

Late-onset depression is common in the elderly but only about 10% of depressed elderly receive treatment. Symptoms of depression include depressed mood, diminished interest or pleasure in activities, significant weight loss or gain, change in sleep patterns, fatigue or loss of energy, diminished ability to think or concentrate, or indecisiveness. Unfortunately, “depression in the elderly is also frequently confused with the effects of multiple illnesses and the medicines used to treat them”

(Goldberg, 2018). Many risk factors for depression exist at all age levels but some appear to be unique to aging, such as:

- Stroke, heart disease, diabetes, cancer, thyroid disorders, and chronic pain
- Parkinson’s and Alzheimer’s disease
- Side effects of medications
- Psychological risk factors such as loneliness, impaired social supports, bereavement, and loss of independence

(Kimmel, 2018; Goldberg, 2018)

In addition to the obvious decline in lifestyle, if left untreated, depression can adversely affect the course of treatment for other chronic diseases. Depressed older adults “are more likely to be admitted to a hospital or a nursing home, visit their physician, visit an emergency room, and more often prescribed multiple medications” (Substance Abuse and Mental Health Services Administration [SAMHSA], 2011, p.8).

Despite the fact that depression is more often diagnosed in women, the highest suicide rates for the elderly are in men, as shown in the following chart:

Deaths by Suicide (2017) (Rate per 100,000)		
Age	Male	Female
65-74	25.9	6.2
75-84	35.8	4.3
85+	50.6	3.5
(CDC, 2019a)		

OTHER ISSUES AFFECTING THE ELDERLY

Elder Abuse

Elder abuse generally refers to intentional actions that cause harm or create a serious risk of harm (whether or not harm is intended) to a vulnerable elder by a caregiver or other person who stands in a trust relationship to the elder whether in the home or an institutional setting (National Center on Elder Abuse [NCEA], n.d.). Elder abuse laws vary from state to state, but most abuse cases fall under the following categories:

- Physical abuse
 - Emotional or psychological abuse
 - Financial or material exploitation
 - Neglect or abandonment by caregivers
 - Sexual Abuse
 - Self-neglect
- (NCEA, n.d.)

Many factors contribute to elder abuse by the caregiver including:

- Alcohol and drug dependency
- Physical and mental stress of caregiving
- Social isolation (elder and caregiver alone together)
- Intensity of the elder’s illness or dementia
- History of violence in family relationships

(Robinson, Saisan, and Segal, 2019)

In 2018, the Texas Department of Adult Protective Services (APS) completed 84,439 investigations of abuse, neglect, or exploitation involving adults, 7,657 of which were in the KCF counties of interest. Of these, 32,878 were confirmed (3,027 in San Antonio) (DFPS, 2019).

Financial Independence

With a longer lifespan comes the challenge of maintaining financial independence. Many seniors struggle to make ends meet. The average income in 2018 for males (living alone) over 65 was \$42,709 and for females (living alone) was \$35,670 (U.S. Census Bureau, 2019a). Major sources of income for seniors in 2018 were reported as social security (84%), income from assets (65%), retirement income (43%) and earnings (24%) (U.S. Census Bureau, 2019b).


Social security benefits in 2018 extended to 67.9 million people (55% women, 45% men) from programs administered by the Social

Security Administration (SSA). Most (62 million) received old-age, survivors, and disability insurance benefits only. 5.4 million received Supplemental Security Income (SSI) only, and 2.7 million received payments from both programs (SSA, 2019, p.32).

“In 2017, 4,681,000 people age 65 and over (9.2%) were below the poverty level” (U.S. Administration on Community Living [ACL], 2018, p.10). Another 2.5 million or 5% of older adults were classified as "near-poor" (income between the poverty level and 125% of this level). Using the U.S. Census Bureau’s Supplemental Poverty Measure (SPM) the poverty level for persons 65 and older was 14.1%.

Older women had a higher poverty rate (10.5%) than older men (7.5%). Older persons living alone were much more likely to be poor (16.7%) than were older persons living with families (5.7%). The highest poverty rates were experienced among older Hispanic women (37.2%) who lived alone (ACL, 2018, p.10).

The following chart represents income reported by seniors in 2017 (ACL, 2018, p.9):



Living Arrangements

The majority (59%) of persons aged 65 and older (16.7 million or 72% of older men, and 13.5 million or 48% of older women) lived with their spouse in 2018. Approximately 28% (4.8 million or 21% of older men, and 9.5 million or 34% of older women) lived alone (ACL, 2018, p.4).

A relatively small number (1.2 million) of the 65+ population in 2017 lived in nursing homes. However, the percentage increased dramatically with age, ranging (in 2017) from

1% for persons 65-74 years to 3% for persons 75-84 years and 9% for persons 85+ (ACL, 2018, p.4).

Although the majority of elderly persons do not require long-term assistance at any given time, most will require assistance at some point in their lives. Independent living programs are an alternative to nursing home placement for many seniors. They are designed to enable seniors to live and thrive in their own homes with the assistance of a network of supportive services. Often referred to as *Retirement Communities* or *Senior Apartments*, independent living communities are generally designed with housing that is friendlier to older adults (easier to access and navigate) and many include more support and recreational options than are readily available to seniors in regular communities (Robinson, Saisan & White, 2019).

Assisted living facilities offer alternatives for seniors that cannot live on their own, but do not need intensive medical or nursing care. While varying in services provided, meals, housing, personal care/support, social activities, security, transportation, and health care management are usually present (Robinson, Saisan & White, 2019).

Nursing homes are the most recognized option of facilities that provide a wide range of long-term care services designed to assist the elderly. Meals, medical, and personal care are all provided, and many provide short-term stays following a hospitalization (Robinson, Saisan & White, 2019).

Continuing Care Retirement Communities (CCRCs) are facilities that offer more than one kind of housing and different levels of care all in the same place to meet the changing needs of the elderly. In the same community, there may be individual homes or apartments for residents who still live on their own, an assisted living facility for people who need some help with daily care, and a nursing home for those who require higher levels of care. While CCRCs provide many advantages, they come with a cost. Nearly two-thirds of these communities charge an average entry fee of \$329,000 with

monthly maintenance fees ranging from \$2,000 to \$4,000 (AARP, 2019).

In 2017, 93% of persons age 65+ were covered by Medicare, which covers mainly acute care services but requires beneficiaries to pay part of the cost, leaving about half of health spending to be covered by other sources (ACL, 2018, p.13).

Caregivers for the elderly

There were approximately 34.2 million individuals caring for someone over the age of 50 in 2015. Caregivers include family, friends and neighbors, with an adult child being the person most likely to be providing care (National Alliance for Caregiving [NAC], 2015).

Caregivers provide assistance to those who need aid in some way. In 2018, 16.2 million family and friends provided 18.5 billion hours of unpaid care to those with Alzheimer's and other dementias (Alzheimer's Association, 2019).

More women than men are caregivers (60% women) and the average age of a female caregiver is 50.3 years old. Most caregivers (59%) are employed and work either full or part-time. Many caregivers (60%) report having to decrease hours, rearrange their schedule, or take unpaid leave in order to meet their caregiving obligations. 17% of caregivers reported fair or poor health as compared to 10% of the general population reporting the same conditions (NAC, 2015).

Agencies that provide support to Texas caregivers include the National Family Caregiver Support Program, Texas Department of Aging and Disability Services (DADS), and Area Agencies on Aging. Working in partnership, these agencies provide five basic services for family caregivers:

- Information about available services
- Assistance in gaining access to supportive services
- Individual counseling, support groups, and caregiver training to assist caregivers in making decisions and solving problems relating to their roles
- Respite services to temporarily relieve

caregivers of their responsibilities

- Supplemental services, on a limited basis, to complement the care provided by caregivers

(ACL, 2019; Texas Health and Human Services [TxHHS], n.d.)

Several centralized sources exist to help seniors find the services that are available:

- *The Eldercare Locator* is a free national service administered by the U.S.

Administration on Aging (AOA) and the National Association of Area Agencies on Aging that can help seniors and their families locate local agencies in every community within the U.S. (AOA, n.d.)

- *Ageing and Disability Resource Center* is a service of Texas Health and Human Services to locate services in Texas (TxHHS, n.d.)

Household Characteristics, 2018												
	United States		Texas		Bexar		Bandera		Comal		Kendall	
		% of total		% of total		% of total		% of total		% of total		% of total
Total Population	322,903,030		27,885,195		1,925,865		21,763		135,097		41,982	
Population 65 years & over	49,238,581	15.2%	3,337,814	12.0%	223,805	11.6%	5,633	25.9%	23,967	17.7%	7,894	18.8%
Total Households	119,730,128		9,553,046		632,574		8,242		48,9031		14,159	
With one or more 65 & older	34,395,751	28.7%	2,316,951	24.3%	153,731	24.3%	3,635	44.1%	15,861	3.2%	4,799	33.9%
Householder 65+ living alone	12,868,890	37.4%	784,778	33.9%	53,266	34.6%	1,041	28.6%	4,663	29.4%	1,341	27.9%

(U.S. Census Bureau, 2020a&c)

REFERENCES

AARP. (2019). *How continuing care retirement communities work*. Retrieved from <https://www.aarp.org/caregiving/basics/info-2017/continuing-care-retirement-communities.html>

Alzheimer’s Association. (2019). *2019 Alzheimer’s disease facts and figures*. Retrieved from <http://www.alz.org/facts/#impact>

America’s Health Rankings. (2018). *Senior report 2018*. Retrieved from <https://assets.americashealthrankings.org/app/uploads/ahrsenior18-finalv1.pdf>

Besdine, R. (2019). *Changes in the body with aging*. Retrieved from <http://www.merckmanuals.com/home/older-people%e2%80%99s-health-issues/the-aging-body/changes-in-the-body-with-aging>

Centers for Disease Control and Prevention [CDC]. (2019a). *Health, United States, 2018: Data finder: Table 009 Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950-2017*. Retrieved from [https://www.cdc.gov/nchs/hus/contents2018.htm?search=Deaths,Older persons](https://www.cdc.gov/nchs/hus/contents2018.htm?search=Deaths,Older%20persons)

Centers for Disease Control and Prevention [CDC]. (2019b). *National health interview survey: Tables of summary health statistics: 2018*. Retrieved from <https://www.cdc.gov/nchs/nhis/SHS/tables.htm>

Denio, A. (2012). *Vitamin D deficiency: the silent epidemic of the elderly*. Retrieved from <http://www.iscd.org/publications/osteoflash/vitamin-d-deficiency-the-silent-epidemic-of-the-elderly/>

Goldberg, J. (2018). *Depression in the elderly*. Retrieved from <https://www.webmd.com/depression/guide/depression-elderly#1>

Goldsmith, T. (2020). *An introduction to biological aging theory (2nd ed. Revision 2)*. Retrieved from http://www.azinet.com/aging/aging_theory_introduction.pdf

Health Resource & Services Administration. (2019). *The “Loneliness epidemic.”* Retrieved from <https://www.hrsa.gov/enevs/past-issues/2019/january-17/loneliness-epidemic>

Holt-Lunstad, J. (2018). *The Potential public health relevance of social isolation and loneliness: Prevalence, epidemiology, and risk factors*. Retrieved from <https://academic.oup.com/ppar/article/27/4/127/4782506>

Kimmel, R. (2018). *Depression- older adults*. Retrieved from <https://medlineplus.gov/ency/article/001521.htm>

Kweder, H. and Eidi, H. (2018). *Vitamin D deficiency in elderly: Risk factors and drugs impact on vitamin D status*. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6178567/>

National Alliance for Caregiving (NAC). (2015). *Caregiving in the U.S. 2015: A Focused look at those caring for someone age 50+*. Retrieved from <https://www.caregiving.org/research/caregivingusa/>

National Center on Elder Abuse [NCEA]. (n.d.). *FAQs: What is elder abuse?* Retrieved February 7, 2020, from <https://ncea.acl.gov/FAQ.aspx>

National Council on Aging [NCOA]. (n.d.a). *5 Facts about malnutrition*. Retrieved February 5, 2020, from <https://www.ncoa.org/healthy-aging/chronic-disease/nutrition-chronic-conditions/why-malnutrition-matters/infographic-facts-malnutrition/>

National Council on Aging [NCOA]. (n.d.b). *SNAP and senior hunger facts*. Retrieved February 5, 2020, from

- <https://www.ncoa.org/news/resources-for-reporters/get-the-facts/senior-hunger-facts/>
- National Institute on Aging (NIA). (n.d.). *Biology of aging*. Retrieved January 27, 2020, from <https://www.nia.nih.gov/about/budget/biology-aging-3>
- National Institute on Aging (NIA). (2018). *Heart health and aging*. Retrieved from <https://www.nia.nih.gov/health/heart-health-and-aging>
- National Institute on Deafness and other Communication Disorders [NIDCD]. (2018). *Age related hearing loss*. Retrieved from <https://www.nidcd.nih.gov/health/age-related-hearing-loss>
- National Institute on Deafness and other Communication Disorders [NIDCD]. (2019). *Quick statistics about taste and smell*. Retrieved from <https://www.nidcd.nih.gov/health/statistics/quick-statistics-taste-smell>
- National Institutes of Health (NIH): Medline Plus. (2020a). *Aging changes in body shape*. Retrieved from <https://www.nlm.nih.gov/medlineplus/ency/article/003998.htm>
- National Institutes of Health (NIH): MedlinePlus. (2020b). *Aging changes in bones, muscles, joints*. Retrieved from <http://www.nlm.nih.gov/medlineplus/ency/article/004015.htm>
- National institutes of Health (NIH): Medline Plus. (2020c). *Aging changes in the nervous system*. Retrieved from <https://www.nlm.nih.gov/medlineplus/ency/article/004023.htm>
- National Institutes of Health (NIH): MedlinePlus. (2020d). *Aging changes in the senses*. Retrieved from <http://www.nlm.nih.gov/medlineplus/ency/article/004013.htm>
- National Institutes of Health [NIH]: MedlinePlus. (2020e). *Aging changes in skin*. Retrieved from <http://www.nlm.nih.gov/medlineplus/ency/article/004014.htm>
- Robinson, L., Saisan, J., and Segal, J. (2019). *Elder abuse and neglect*. Retrieved from <https://www.helpguide.org/articles/abuse/elder-abuse-and-neglect.htm>
- Robinson, L., Saisan, J., and White, M. (2019). *Senior housing options*. Retrieved from <https://www.helpguide.org/articles/alzheimers-dementia-aging/senior-housing.htm>
- Ruiz, A. (2019). *Effects of aging on the digestive system*. Retrieved from <http://www.merckmanuals.com/home/digestive-disorders/biology-of-the-digestive-system/effects-of-aging-on-the-digestive-system#v752160>
- Sebastian, R., Cleveland, L., Goldman, J., and Moshfegh, A. (2007). *Older adults who use vitamin/mineral supplements differ from nonusers in nutrient intake adequacy and dietary attitudes*. Retrieved from <http://hdl.handle.net/10113/23063>
- Substance Abuse and Mental Health Services Administration (SAMHSA). (2011). *Depression and older adults: Key issues*. Retrieved from <https://store.samhsa.gov/system/files/sma11-4631-keyissues.pdf>
- Texas Demographic Center [TDC]. (2019). *2018 Estimated population of Texas, its counties, and places*. Retrieved from <https://demographics.texas.gov/>
- Texas Department of Family and Protective Services [DFPS]. (2019). *Adult Protective Services: DFPS data book*. Retrieved from https://www.dfps.state.tx.us/About_DFPS/Data_Book/Adult_Protective_Services/Investigations/Findings.asp
- Texas Health and Human Services (TxHHS). (n.d.). *Aging and disability resource center*. Retrieved February 14, 2020, from <https://hhs.texas.gov/services/aging/long-term-care/aging-disability-resource-center>
- U.S. Administration on Aging [AOA]. (n.d.). *Welcome to Eldercare locator*. Retrieved February 14, 2020, from <https://eldercare.acl.gov/Public/Index.aspx>
- U.S. Administration on Community Living [ACL]. (2018). *Profile of older Americans: 2018*. Retrieved from <https://acl.gov/sites/default/files/Aging%20and%20Disability%20in%20America/2018OlderAmericansProfile.pdf>
- U.S. Administration on Community Living [ACL]. (2019). *National family caregiver support program*. Retrieved from <https://www.acl.gov/programs/support-caregivers/national-family-caregiver-support-program>
- U.S. Census Bureau. (2019a). *2019 CPS Household income table: HINC-02. Age of householder-households, by total money income in 2018, type of household, race and Hispanic origin of householder*. Retrieved from https://www.census.gov/data/tables/2019/demo/cps/hinc-02.html#par_textimage_28
- U.S. Census Bureau. (2019b). *PINC-08. Source of income-people 15 years old and over, by income of specified type, age, race, Hispanic origin, and sex*. Retrieved from <https://www.census.gov/data/tables/time-series/demo/income-poverty/cps-pinc/pinc-08.html>
- U.S. Census Bureau. (2020a). *ACS demographic and housing estimates: 2018 ACS 5-year estimates data profiles: Table DP05*. Retrieved from https://data.census.gov/cedsci/table?q=Population%20Total&g=010000US_0400000US48_0500000US48019,48029,48091,48259&hidePreview=true&t=Population%20Total&table=DP05&tid=ACSDPY2018.DP05&lastDisplayedRow=93&vintage=2018&moe=false
- U.S. Census Bureau. (2020b). *Disability characteristics: 2018 ACS 1-year estimates subject tables: Table S1810*. Retrieved from <https://data.census.gov/cedsci/table?q=S1810%3A%20DISABILITY%20CHARACTERISTICS&hidePreview=true&tid=ACSST1Y2018.S1810&vintage=2018>
- U.S. Census Bureau. (2020c). *Households by presence of people 65 years and over, household size and household type: 2018: ACS 5-year estimates detailed tables: Table B11007*. Retrieved from https://data.census.gov/cedsci/table?q=table%20B11007&g=0100000US_0400000US48_0500000US48029,48091,48019,48259&hidePreview=true&tid=ACSDT5Y2018.B11007&vintage=2018&moe=false
- U.S. Census Bureau. (2020d). *Population 65 Years and over in the United States: 2018: ACS 5-year estimates subject tables: Table S0103*. Retrieved from <https://data.census.gov/cedsci/table?q=Table%20PEPAGESEX&g=&table=S0103&tid=ACSST5Y2010.S0103&vintage=2018&t=Age%20and%20Sex%3AOlder%20Population&lastDisplayedRow=19&mode=>
- U.S. Census Bureau. (2020e). *Projected age groups and sex composition of the population: Projections for the United States: 2017-2060: Table 2*. Retrieved from <https://www.census.gov/data/tables/2017/demo/popproj/2017-summary-tables.html>
- U.S. Department of Health and Human Services [HHS]. (2016a). *The Challenge of managing multiple chronic conditions*. Retrieved from <https://www.hhs.gov/ash/about-ash/multiple-chronic-conditions/about-mcc/the-challenge-of-managing-multiple-chronic-conditions/index.html>
- U.S. Social Security Administration (SSA). (2019). *Fast facts and figures about social security*. Retrieved from https://www.ssa.gov/policy/docs/chartbooks/fast_facts/2019/fast_facts19.pdf
- World Health Organization [WHO]. (2017). *Mental health of older adults*. Retrieved from <http://www.who.int/mediacentre/factsheets/fs381/en/>