

Alzheimer's disease is the most common form of dementia, constituting 60-80% of all dementia cases. The fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) by the American Psychiatric Association includes two dementia categories in the diagnostic classification of neurocognitive disorders:

- *Major neurocognitive disorder* classification requires that an "individual must have evidence of significant cognitive decline, and the decline must interfere with independence in everyday activities (for example, assistance may be needed with complex activities such as paying bills or managing medications)"
- *Mild neurocognitive disorder* classification requires that "individual must have evidence of modest cognitive decline, but the decline does not interfere with everyday activities (individuals can still perform complex activities such as paying bills or managing medications, but the activities require greater mental effort)"
(Alzheimer's Association, n.d.)

Alzheimer's disease is a degenerative brain disease where nerve cells in the brain are damaged and no longer function properly (Alzheimer's Association, 2022a). Plaques represent a type of physical abnormality seen in brains of people with Alzheimer's disease. Plaques occur when the protein amyloid-beta builds up between neurons, causing the neurons to clump and die. Another brain abnormality in Alzheimer's patients is the presence of tangles.

Tangles occur when the protein tau causes neurons to twist, resulting in tremendous neuron damage (Mayo Clinic, 2022).

ALZHEIMER'S CONTINUUM

"The progression of Alzheimer's disease from brain changes that are unnoticeable to the person affected to brain changes that cause problems with memory and eventually physical disability is called the Alzheimer's disease continuum" (Alzheimer's Association, 2022a, p9).

1. *Preclinical Alzheimer's disease*. In this stage, there are no symptoms but there are measureable changes in the brain, cerebrospinal fluid and/or blood evidenced by biomarkers. It is estimated that brain changes occur as many as 20 years before symptoms are evident.
2. *Mild Cognitive Impairment (MCI) due to Alzheimer's disease*. Includes individuals who have mild but measureable changes in thinking abilities that do not affect the ability to carry out everyday activities. It is estimated that 15-20 percent of people age 65 and older have MCI.
3. *Dementia due to Alzheimer's disease*. Cognitive impairment interferes with an individual's ability to function in daily life and is caused by Alzheimer's disease-related changes in the brain. This phase is further broken down into three stages:
 - a. Mild – symptoms interfere with some everyday activities but may still be able to drive, work, and participate in favorite activities
 - b. Moderate – "individuals experience more problems with memory and

language, and are more likely to become confused, and find it harder to complete multistep tasks such as bathing and dressing. They may become incontinent at times, and they may start having personality and behavioral changes”

- c. Severe – individuals need assistance with daily living activities and around-the-clock care

(Alzheimer's Association, 2022a, p.10)

THE PROGRESSION OF ALZHEIMER'S DISEASE

Alzheimer's disease progressively affects memory, thinking, and behavior. The *Global Deterioration Scale*, also called the *Reisberg Scale*, divides the progression of Alzheimer's disease into seven key stages of development:

Stage One: No cognitive impairment

- No memory problems evident

Stage Two: Very mild cognitive decline

- Memory lapses for familiar words and names
- Lapses not evident to friends, family, co-workers, or doctors

Stage Three: Mild cognitive decline

- Friends, family, and co-workers begin to notice deficiencies in memory
- Problems with memory, retention, and concentration
- Difficulty naming objects or people
- Decreased performance of daily activities
- Decline in ability to plan or organize

Stage Four: Moderate cognitive decline

- Clear cut deficiencies in knowledge of recent events
- Decreased capacity to perform complex tasks such as planning Thanksgiving dinner or managing finances
- Demeanor may be subdued or withdrawn

Stage Five: Moderately severe cognitive decline

- Major gaps in memory and cognitive functioning
- Need for assistance with daily functions
- Confusion about location and date

Stage Six: Severe cognitive decline

- Increased memory deficits

- Significant personality changes
- Difficulty recalling personal history
- Tendency to wander and become lost
- Impaired ability to execute common actions such as dressing oneself

Stage Seven: Very severe cognitive decline

- Little to no ability to speak
- Failure to recognize objects and people
- Loss of voluntary muscle control
- Assistance required for all daily functions

(Alzheimer Society Canada, 2016)

Patients may not exhibit all signs and symptoms of each stage, nor does the disease progress at an equal rate in every patient. The Alzheimer's Association reports “studies indicate that people age 65 and older survive an average of 4 to 8 years after a diagnosis of Alzheimer's disease, yet some live as long as 20 years with Alzheimer's dementia” (Alzheimer's Association, 2022a, p.35).

PREVALENCE AND RISK FACTORS

Currently, an estimated 6.5 million people in the United States are living with Alzheimer's disease. The majority, 6.47 million, are over age 65, while only 200,000 are younger than 65. 72.6% of seniors with Alzheimer's disease are age 75 or older (Alzheimer's Association, 2022a, p.19).

The single greatest known risk factor for Alzheimer's disease is age. Approximately 11% of Americans over the age of 65 have Alzheimer's disease with higher incidence as age increases. Approximately 33% of seniors 85 and older have Alzheimer's disease (Alzheimer's Association, 2022a).

It is estimated that Alzheimer's disease effects approximately one in ten people aged 65 and older. As a result of a longer life expectancy, women are more likely to develop Alzheimer's disease than men and currently two-thirds of Alzheimer's patients are women. Alzheimer's disease and dementias account for approximately 12 percent of women as opposed to only 9 percent of men over age 65. Of this age group, women represent 4 million of Americans with Alzheimer's, men represent 2.5 million but there is no indication that women were more likely than men to develop the

disease at any given age (Alzheimer’s Association, 2022a).

“In 2011, the largest ever demographic generation of the American population — the baby boom generation — started reaching age 65, the age at which the risk for Alzheimer’s and other dementias begins to increase. By 2030, the segment of the U.S. population age 65 and older will have grown substantially, and the projected 74 million older Americans will make up over 20% of the total population (up from 18% in 2022” (Alzheimer’s Association, 2022a, p.27). As Americans live longer and the baby boomer generation ages, rates of Alzheimer’s disease are also projected to grow exponentially:

YEAR	PROJECTED # OF INDIVIDUALS AGE 65 AND OVER WITH ALZHEIMER’S DISEASE (IN MILLIONS)
2020	6.1m
2030	8.5m
2040	11.2m
2050	13.8m

(Alzheimer’s Association, 2022a, p. 26)

Other risk factors for Alzheimer’s disease include family history, genetics, environmental factors, and history of severe head injury. It is believed that certain genetic factors may increase the potential for developing Alzheimer’s disease. For example, the genetic presence of particular forms of the protein Apolipoprotein-E (ApoE4) gene, which occurs in people with late-onset Alzheimer’s disease, has been identified as a potential risk factor. Early-onset Alzheimer’s disease has been linked to mutations in three genes. “A blood test can identify which APOE alleles a person has, but results cannot predict who will or will not develop Alzheimer’s disease” (National Institute on Aging, 2019). Except for research (clinical trials) and to help doctors diagnose early-onset Alzheimer’s, genetic testing is not recommended. Other risk-factor genes are under investigation.

Researchers are also investigating the association between various lifestyle factors related to the cardiovascular system and their influence on Alzheimer’s disease. Because the brain contains numerous blood vessels,

conditions such as smoking, obesity (especially in midlife), diabetes, high cholesterol in midlife, and hypertension in midlife may be linked to the onset of Alzheimer’s disease (Alzheimer’s Association, 2022a, p. 82).

Recent research suggests that the prevalence and risk of Alzheimer’s disease is higher in both Hispanics and African Americans, as compared with other Americans, perhaps due to socioeconomic factors as well as higher rates of cardiovascular diseases. Specific prevalence rates among each racial and ethnic group remain unclear. This ambiguity is largely attributable to African Americans and Hispanic Americans avoiding or delaying treatment, causing the disease to go undiagnosed (Alzheimer’s Association, 2022a).

“Most people with Down syndrome develop Alzheimer’s. This may be because people with Down syndrome have an extra copy of chromosome 21, which contains the gene that generates harmful amyloid” (National Institute on Aging, 2021).

Additionally, new research has linked fewer years of education to an increased risk of developing Alzheimer’s disease. The exact cause of this association remains unclear. Some researchers theorize that additional years of education allow for the development of more brain synapses resulting in a greater reserve as individuals age while others believe that those with lower education also are subject to other risk factors common to lower socioeconomic groups including increased risk for disease in general and less access to medical care (Alzheimer’s Association, 2022a).

TREATMENT

No known cure for Alzheimer’s disease exists, but there are two types of FDA approved drugs which can delay its progression:

1. *Cholinesterase inhibitors*
 - a. Prevent the breakdown of the chemical acetylcholine (which aids memory and thinking)
 - b. Delay symptoms for 6-12 months in approximately half of patients.

- c. Used for those with mild to moderate Alzheimer's disease
2. N-methyl D-aspartate (NMDA) antagonists
 - a. Help regulate the activity of glutamate (which aids information processing)
 - b. Used for those with moderate to severe Alzheimer's disease

“Because NMDA antagonists work very differently from cholinesterase inhibitors, the two types of drugs can be prescribed in combination” (National Institute on Aging, 2018, p.3).

Several drugs targeting beta-amyloid, the chief component of plaques, are under investigation. “While scientists aren't sure what causes cell death and tissue loss during the course of Alzheimer's, amyloid plaques are one of the potential contributors, and preventing beta-amyloid buildup may provide benefit (Alzheimer's Association, 2022e).

Other medical treatments seek to alleviate Alzheimer's symptoms beyond cognitive impairment such as sleeplessness, agitation, wandering, anxiety, aggression, and depression (National Institute on Aging, 2021).

RESPIRE SERVICES

Proper care and support is essential to reduce the complications common to Alzheimer's, including pneumonia, infections, falls, fractures, and malnutrition (Mayo Clinic, 2022). Currently, over 16 million Americans, usually relatives or friends, provide unpaid care for individuals suffering from dementia or Alzheimer's disease. “In 2021, caregivers of people with Alzheimer's or other dementias provided an estimated 16 billion hours of informal (that is, unpaid) assistance, a contribution to the nation valued at \$271.6 billion” (Alzheimer's Association, 2022a, p38). In Texas, 1.1 million caregivers provided over 1.8 billion hours of unpaid care in 2021.

Because it is possible for a patient to live with the disease for many years, the emotional and financial burden on caregivers can be overwhelming. Consequently, it is common for the caregivers themselves to suffer from illnesses such as heart disease, high-blood pressure, and poor mental health (depression).

These health issues are often directly related to the stress and anxiety that comes with caring for a loved one with Alzheimer's disease (Alzheimer's Association, 2022a). Fortunately, there are several types of services that exist to support caregivers.

One option available to caregivers is in-home care services, which involves a professional coming to the home to assist the caregiver and individual suffering from dementia with a variety of tasks. The following are common in-home care services:

- **Companion services:** assist caregiver with supervision, recreational activities, and visiting
- **Personal care services:** provide help with bathing, dressing, toileting, exercising, or other personal care needs
- **Homemaker services:** provide assistance with housekeeping, shopping, and meal preparation
- **Skilled care:** assists with wound care, injections, physical therapy and other medical needs, usually conducted by a licensed health professional

(Alzheimer's Association, 2022c)

Outside of the home, adult day services provide those suffering from dementia or Alzheimer's disease opportunities for social interaction and a safe environment to participate in activities. These adult day programs provide a variety of services, including meals, social and educational activities, exercise, various types of therapy, counseling, and medical treatment. Typically, these services are available daily and most programs are flexible with attendance requirements. “Over 80% of participants attend full days and 46% attend five days per week, enabling family caregivers to remain in the workforce” (National Adult Day Services Association, n.d.). Most adult day centers also provide a variety of caregiver support programs including education, support groups, and counseling.

Locating respite services for adults can be challenging for caregivers. There are several online locator services that can help. The Texas Health and Human Services (HHS) website lists

51 respite care providers in Bexar County. (HHS, n.d.).

Long-term respite care facilities include assisted living facilities, residential care facilities, and nursing homes. These programs provide overnight, weekend, or long-term stays when a caregiver experiences illness or other emergency situations. Services include meals, daily living help, therapeutic activities, and a safe setting to reduce wandering (Alzheimer's Association, 2022d).

For those individuals who need specialty care, there are several different kinds of facilities which assist in care for different stages of Alzheimer's. These facilities include:

- **Retirement housing** for early stage Alzheimer's patients who are still able to care for themselves and require no more than limited supervision, social activities, and transportation
- **Assisted living** (also called *board and care*, *adult living* or *supported care*) offers moderately independent Alzheimer's patients housing, meals, supportive services, and health care (these facilities are not government regulated)
- **Nursing homes** (also called *skilled nursing facility*, *long-term care facility* or *custodial care*) offer around-the-clock care and long-term medical treatment for mid to late-stage Alzheimer's patients (these facilities are regulated by both state and federal governments)
- **Alzheimer's special care units (SCUs)** (also called *memory care units*) assist those with dementia or Alzheimer's disease and can be found in many types of care facilities
- **Continuing care retirement communities (CCRC)** give either independent, assisted living, or nursing home levels of care depending on individual need

(Alzheimer's Association, 2022d)

THE HIGH COST OF ALZHEIMER'S DISEASE

Alzheimer's disease also creates an enormous economic responsibility for caregivers, the government, through Medicare and Medicaid, and independent businesses. The following

statistics, provided by the Alzheimer's Association (2022a, p. 69-70), represent typical costs associated with Alzheimer's care.

Average costs for Alzheimer's services:

- **Adult day care service** – \$75 per day
- **Assisted living** – approximately \$4,429 per month or \$53,148 per year (dementia care often results in additional charges)
- **Nursing Homes** – about \$299 per day or \$109,135 per year (for a semi-private room the cost is around \$263 per day or \$95,995 per year)
- **Home care** – cost for a nonmedical home health aide was \$24 per hour or \$4,652 per month

The high costs incurred from Alzheimer's disease are also reflected on United States businesses due to caregiver absenteeism, loss of productivity, and employee turnover. Independent businesses accrue additional costs from health care, long-term care, and hospice for people with Alzheimer's disease and other dementias. The total estimated cost for health care and long-term care for people aged 65 and older with Alzheimer's disease and other types of dementia totaled \$321 billion for the year 2021. By the year 2050, the cost is expected to rise to just under \$1 trillion (Alzheimer's Association, 2022a, p. 79).

MORTALITY

Through advancements in research and technology, there are ways to diagnose Alzheimer's such as blood tests that measure beta-amyloid levels. The disease used to be definitively diagnosed through an autopsy (National Institute on Aging, 2021). In any given year, one-third of all seniors who die are found to have Alzheimer's or some form of dementia. For people aged 65 and older, Alzheimer's disease is the fifth-leading cause of death in the United States, and the sixth-leading cause of death across all age groups (Alzheimer's Association, 2022a).

For the year 2019, Alzheimer's disease was the underlying cause of death for 10,101 individuals in Texas. The U.S. annual Alzheimer's death rates show that deaths

attributed to the disease increase drastically with age. The increase in deaths attributed to Alzheimer's disease over time has disproportionately affected the oldest populations: "Between 2000 and 2019, the death rate from Alzheimer's increased 33% for people aged 65 to 74, but increased 51% for people age 75 to 84 and 78% for people age 85 and older" (Alzheimer's Association, 2022a, p.34).

"Scientists have developed methods to measure and compare the burden of different diseases on a population in a way that takes into account not only the number of people with the condition, but also the number of years of life lost due to that disease and the number of healthy years of life lost by virtue of being in a state of disability. One measure of disease burden is called disability-adjusted life years (DALYs), which is the sum of the number of years of life lost (YLLs) due to premature mortality and the number of years lived with disabilities (YLDs), totaled across all those with the disease or injury...Alzheimer's disease rose from the 12th most burdensome disease or injury in the United States in 1990 to the sixth in 2016 in terms of DALYs. In 2016, Alzheimer's disease was the fourth highest disease or injury in terms of YLLs and the 19th in terms of YLDs" (Alzheimer's Association, 2022a, p35).

COVID-19 CONSIDERATIONS

Services and programs for Alzheimer's Dementia patients have been impacted by the COVID-19 Pandemic. Patients with Alzheimer's, while not at an increased risk to contract the virus, can find themselves at higher risk of being impacted by it. Through things like forgetting to wash hands and take necessary precautions, people with Alzheimer's may be at higher risk. Similarly, "COVID-19 may worsen cognitive impairment due to dementia (Alzheimer's Association, 2022e). Local organizations like the Alzheimer's Association in San Antonio are offering virtual services and services over the phone as well as other safety precautions on site (Alzheimer's Association, 2022e).

REFERENCES

- Alzheimer's Association. (n.d.). *In brief for healthcare professionals: Differentiating dementias*. Retrieved April 27, 2022, from <https://www.alz.org/media/documents/inbrief-differentiating-dementias.pdf>
- Alzheimer's Association. (2022a). *2020 Alzheimer's disease facts and figures*. Retrieved from <https://www.alz.org/media/Documents/alzheimers-facts-and-figures.pdf>
- Alzheimer's Association. (2022b). *In-home care*. Retrieved from <https://www.alz.org/help-support/caregiving/care-options/in-home-care>
- Alzheimer's Association. (2022c). *Residential Care*. Retrieved from <https://www.alz.org/care/alzheimers-dementia-residential-facilities.asp>
- Alzheimer's Association. (2022d). *Treatment horizon*. Retrieved from http://www.alz.org/research/science/alzheimers_treatment_horizon.asp
- Alzheimer's Association. (2022e). *COVID-19, Alzheimer's and dementia: what you need to know*. Retrieved from <https://www.alz.org/alzheimers-dementia/coronavirus-covid-19>
- Alzheimer Society Canada (2016). *Progression*. Retrieved from <https://alzheimer.ca/sites/default/files/documents/Progression-overview--Alzheimer-Society.pdf>
- Mayo Clinic. (2022). *Alzheimer's disease: causes*. Retrieved from <http://www.mayoclinic.org/diseases-conditions/alzheimers-disease/symptoms-causes/dxc-20167103>
- National Adult Day Services Association. (n.d.). *About adult day services*. Retrieved April 27, 2022, from <http://www.nadsa.org/learn-more/about-adult-day-services/>
- National Institute on Aging. (2018). *Alzheimer's disease medications fact sheet*. Retrieved from <https://order.nia.nih.gov/sites/default/files/2018-03/alzheimers-disease-medications-fact-sheet.pdf>
- National Institute on Aging. (2019). *Alzheimer's disease genetics fact sheet*. Retrieved from <https://www.nia.nih.gov/alzheimers/publication/alzheimers-disease-genetics-fact-sheet>
- National Institute on Aging. (2021a). *Alzheimer's disease fact sheet*. Retrieved from <https://www.nia.nih.gov/health/alzheimers-disease-fact-sheet#causes>
- National Institute on Aging. (2021b). *How is Alzheimer's disease diagnosed?* Retrieved from <https://www.nia.nih.gov/health/how-alzheimers-disease-diagnosed>
- Texas Health and Human Services (HHS). (n.d.). *Respite provider search*. Retrieved May 2, 2022, from <https://apps.hhs.texas.gov/taketimetexas/search.cfm>